

Hypertension

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Learn and Live SM

Context Dependency of Serum and Urinary Lithium: Implications for Measurement of Proximal Sodium Reabsorption

Murielle Bochud, Jan A. Staessen, Angela Woodiwiss, Gavin Norton, Marc Maillard and Michel Burnier

Hypertension 2007;49;34-; originally published online Mar 26, 2007;

DOI: 10.1161/HYPERTENSIONAHA.107.089862

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 72514

Copyright © 2007 American Heart Association. All rights reserved. Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hyper.ahajournals.org/cgi/content/full/49/5/e34>

Subscriptions: Information about subscribing to Hypertension is online at
<http://hyper.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax: 410-528-8550. E-mail:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/reprints>

Letter to the Editor

Letters to the Editor will be published, if suitable, as space permits. They should not exceed 1000 words (typed, double-spaced) in length and may be subject to editing or abridgment.

Context Dependency of Serum and Urinary Lithium: Implications for Measurement of Proximal Sodium Reabsorption

To the Editor:

As reported previously in this journal, the endogenous lithium clearance is a marker of proximal tubular sodium handling.^{1,2} To investigate the context dependency of the endogenous lithium clearance, we measured by electrothermal atomic absorption spectrophotometry the lithium concentrations in serum and in exactly timed urine samples in 745 whites (51.5% women) and 266 blacks (62.8%) who were recruited randomly from the population in Belgium¹ and South Africa.³ Mean age (\pm SD) was 40.6 ± 15.8 and 42.6 ± 18.2 years, respectively. The average (\pm SD) concentration of lithium in tap water (1.37 ± 0.56 versus 0.03 ± 0.01 $\mu\text{mol/L}$) and the 24-hour urinary lithium excretion (8.2 ± 5.6 versus 3.1 ± 4.1 μmol per 24 hours) were higher and more dispersed in Belgium than in South Africa ($P < 0.0001$ for differences in means and variances; Figure). These results strongly suggest that the environment, probably via the food chain,⁴ determines the dietary intake of lithium. By contrast, the serum lithium distributions (0.31 ± 0.16 versus 0.32 ± 0.21 $\mu\text{mol/L}$) were very similar in means ($P = 0.58$) but not in variances ($P < 0.0001$). These observations suggest that serum lithium is tightly regulated despite large variations in dietary intake. The rank correlations (\pm SE) between the lithium and creatinine clearances (0.44 ± 0.04 versus 0.42 ± 0.06 ; $P = 0.70$ for between-country difference) and between the serum and urinary lithium concentrations (0.30 ± 0.04 versus 0.22 ± 0.06 ; $P = 0.26$) were similar across countries. The latter

observation suggests that the endogenous lithium clearance, whether expressed as clearance or as fractional excretion, remains a useful marker of proximal sodium reabsorption under various environmental conditions.

Sources of Funding

The study was financed in part by InGenious HyperCare (LSHM-CT-2006-037093), a grant of the European Union.

Disclosures

None.

Murielle Bochud

Community Prevention Unit
University Institute for Social
and Preventive Medicine
Lausanne, Switzerland


Jan A. Staessen

Department of Cardiovascular Diseases
University of Leuven
Leuven, Belgium

Angela Woodiwiss

Gavin Norton

Cardiovascular Pathophysiology
and Genomics Research Unit
University of the Witwatersrand
Johannesburg, South Africa



Distributions of serum and urinary lithium. Outlier values were removed.

Marc Maillard**Michel Burnier***Nephrology Division**University Hospital of Lausanne**Lausanne, Switzerland*

1. Seidlerova J, Staessen JA, Maillard M, Nawrot T, Zhang H, Bochud M, Kuznetsova T, Richart T, Van Bortel LM, Struijker-Boudier HA, Manunta P, Burnier M, Fagard R, Filipovsky J. Association between arterial properties and renal sodium handling in a general population. *Hypertension*. 2006;48:609–615.
2. Eap CB, Bochud M, Elston RC, Bovet P, Maillard M, Nussberger J, Schild L, Shamlaye C, Burnier M. *CYP3A5* and *ABCB1* genes influence blood pressure and response to treatment, and their effect is modified by salt. *Hypertension*. 2007;49:1007–1014.
3. Shiburi CP, Staessen JA, Maseko M, Wojciechowska W, Thijs L, Van Bortel LM, Woodiwiss AJ, Norton GR. Reference values for SphygmoCor measurements in South Africans of African ancestry. *Am J Hypertens*. 2006;19:40–46.
4. Schrauzer GN. Lithium: occurrence, dietary intakes, nutritional essentiality. *J Am Coll Nutr*. 2002;21:14–21.